

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Independents Own

Fenbender 100

Oral Drench for Cattle & Horses

**ACTIVE CONSTITUENT: 100 g/L
FENBENDAZOLE**

For the control of benzimidazole-sensitive mature and immature Roundworms (including inhibited *Ostertagia*), Lungworms and as an aid in the control of Tapeworms in Cattle. For the treatment and control of Roundworms, Bloodworms, Redworms and Pinworms in horses.

INDEPENDENTS OWN FENBENDER 100 ORAL DRENCH FOR CATTLE AND HORSES contains Fenbendazole, a member of the benzimidazole family of chemicals.

It is effective against sensitive strains of the following internal parasites:

CATTLE: For the control of Benzimidazole-Sensitive mature and immature Roundworms (including inhibited *Ostertagia*), Lungworms and as an aid in the control of Tapeworms in cattle:

(*Haemonchus placei*) Barber's pole worm; (*Trichostrongylus axei*) Stomach hair worm; (*Trichostrongylus spp*) Intestinal hair worm or Black scour

worm; (*Bunostomum phlebotomum*) Hookworm; (*Cooperia spp*) Small intestinal worm; (*Nematodirus spp*) Thin necked intestinal worm; (*Oesophagostomum radiatum*) Nodule worm; (*Dictyocaulus viviparus*) Lungworm, (*Ostertagia ostertagi*) Small brown stomach worm including 80% - 90% activity against Hypobiotic larvae (*Moniezia spp*) Tapeworm.

HORSES: For treatment and control of mature and immature Roundworms, Bloodworms, Redworms and Pinworms in horses: (*Strongylus spp*) Large strongyle, (*Cyathostomum spp*, *Triodontophorus spp.*) Small strongyle, (*Parascaris equorum*) Ascarids, (*Oxyuris equi*) Pinworms.

Resistance may develop to any chemical.

DIRECTIONS FOR USE:

SHAKE WELL BEFORE USE.

NOT FOR USE IN GOATS.

This product can be administered concurrently with vaccines, pour-on lice treatments and dipping operations.

CATTLE: For the control of Benzimidazole-Sensitive mature and immature Roundworms (including inhibited *Ostertagia*), Lungworms and as an aid in the control of Tapeworms in cattle:

(*Haemonchus placei*) Barber's pole worm; (*Trichostrongylus axei*) Stomach hair worm; (*Trichostrongylus spp*) Intestinal hair worm or Black scour worm; (*Bunostomum phlebotomum*) Hookworm; (*Cooperia spp*) Small intestinal worm; (*Nematodirus spp*) Thin necked intestinal worm; (*Oesophagostomum radiatum*) Nodule worm; (*Dictyocaulus viviparus*) Lungworm (*Ostertagia ostertagi*) Small brown stomach worm including 80% - 90% activity against Hypobiotic larvae (*Moniezia spp*) Tapeworm.

HORSES: For treatment and control of mature and immature Roundworms, Bloodworms, Redworms and Pinworms in horses:

(*Strongylus spp*) Large strongyle, (*Cyathostomum spp*, *Triodontophorus spp.*) Small strongyle, (*Parascaris equorum*) Ascarids, (*Oxyuris equi*) Pinworms.

Resistance may develop to any chemical

DOSAGE & ADMINISTRATION:

SHAKE WELL BEFORE USE.

CATTLE: 7.5 mg/kg Fenbendazole or 3 mL per 40 kg bodyweight. Cattle in excess of 680 kg to be dosed at 3 mL per 40 kg bodyweight.

| Bodyweight | Dose | No. of Animals 1L Pack |
|--------------|-------|------------------------|
| Up to 80 kg | 6 mL | 167 |
| 81 - 120 kg | 9 mL | 133 |
| 121 - 160 kg | 12 mL | 83 |
| 161 - 200 kg | 15 mL | 67 |
| 201 - 240 kg | 18 mL | 55 |
| 241 - 280 kg | 21 mL | 47 |
| 281 - 320 kg | 24 mL | 41 |
| 321 - 360 kg | 27 mL | 37 |
| 361 - 400 kg | 30 mL | 33 |
| 401 - 440 kg | 33 mL | 30 |
| 441 - 480 kg | 36 mL | 27 |
| 481 - 520 kg | 39 mL | 25 |
| 521 - 560 kg | 42 mL | 23 |
| 561 - 600 kg | 45 mL | 22 |
| 601 - 640 kg | 48 mL | 21 |
| 641 - 680 kg | 51 mL | 19 |

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Check accuracy of drenching gun before and during use. Rinse drenching guns after use.

HORSES: 10 mg/kg Fenbendazole or 5 mL per 50 kg body weight. For the control of *Strongyloides westeri* in foals: 50 mg/kg or 5 mL/10 kg body weight. Independents Own Fenbender 100 Oral Drench may be administered to horses by nasal tube, drenching gun or dosage syringe.

Worm parasites may develop resistance to Fenbendazole. If this is suspected, please contact your local veterinary practitioner or advise the manufacturer.

WITHHOLDING PERIODS:

CATTLE: MEAT: DO NOT USE less than 21 days before slaughter for human consumption.

MILK: Zero (0) Days

HORSES: DO NOT USE less than 28 days before slaughter for human consumption.

TRADE ADVICE – Export Slaughter Interval:

CATTLE: DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of approval. Before using this product confirm the ESI from Apparent Pty Ltd on 0411 227 338 or the APVMA website (www.apvma.gov.au/residues/)

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE: Store below 30°C (Room temperature) with top secure - Avoid Freezing.

SAFETY DIRECTIONS: Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET: Additional information is listed in the Safety Data Sheet.

APPARENT PTY LTD

Suite G.08, 762 Toorak Road, Hawthorn East, Vic. 3123,
PO Box 3092 Cotham PO
KEW VIC 3101

P 03 9822 1321 **M** 0411 227 338

E wardell@bigpond.net.au

APVMA Approval Number:
81050/102239